



**The Use of Biotelemetry in Controlling
Carp (*Cyprinus carpio*) in Lakes
Crescent and Sorell by the Inland
Fisheries Service Carp Management
Program.**

Technical Report No 1, June 2003

Prepared by Alasdair Macdonald.
Technical Officer - Carp Management Program.

Inland Fisheries Service 2003
Tasmania
Australia

© Inland Fisheries Service, July 2003

© For further information contact:

Inland Fisheries Service
PO Box 288
Moonah Tas 7009
Telephone: (03) 6233 4140
Facsimile: (03) 6233 4141
Email: infish@ifs.tas.gov.au

Find further information about IFS on the internet site <http://www.ifs.tas.gov.au>

This report is part of a series of documents, which provide information and details of carp eradication efforts in lakes Sorell and Crescent as part of the Lakes Sorell and Crescent Carp Management Project.

The aim of the project is to control the spread of carp within the state of Tasmania, with a view to their eradication.

Citation: Macdonald A. (2003). The Use of Biotelemetry in Controlling Carp (*Cyprinus carpio*) in lakes Crescent and Sorell by the Inland Fisheries Service Carp Management Program. Technical Report Inland Fisheries Service, Hobart.

Copyright

All material published in this report by the Inland Fisheries Service, as an agent of the Crown, is protected by the provisions of the *Copyright Act 1968* (Cth). Other than in accordance with the provisions of the Act, or as otherwise expressly provided, a person must not reproduce, store in a retrieval system, or transmit any such material without first obtaining the written permission of the Inland Fisheries Service.

Disclaimer

Whilst the Service makes every attempt to ensure the accuracy and reliability of information published in this report, it should not be relied upon as a substitute for formal advice from the originating bodies or Departments. The IFS, its employees and other agents of the Crown will not be responsible for any loss, however arising, from the use of, or reliance on this information.

Contents

1.0 Introduction.....	1
2.0 Telemetry Equipment.....	2
2.1 Transmitters	2
2.2 Telemetry Receiver Systems	5
2.3 Receiving Transmitter Signals	7
3.0 Radio Tracking Techniques	9
4.0 Surgical Implant Techniques.....	12
4.1 Introduction.....	12
4.2 Surgical Equipment	13
4.3 Sterilisation and fish handling procedures ..	14
4.4 Anaesthesia.....	15
4.5 Surgical Procedure.....	15
5.0 Discussion	19
6.0 References	21

1.0 Introduction

Biotelemetry provides a means to monitor the location, behaviour and physiology of animals in uncontrolled environments, both terrestrial and aquatic. Biotelemetry or 'radio tracking' involves the transfer of information in the form of radio signals from radio transmitters, (which are attached or implanted into animals), to a remote receiver system.

Since 1997, the Carp Management Program, under the Tasmanian Inland Fisheries Service (IFS) has employed the use of biotelemetry technology to identify the movements of carp (*Cyprinus carpio*) in lakes Crescent and Sorell, Tasmania.

Carp were discovered in these lakes in 1995 and since then considerable resources have been allocated to reduce their abundance and maintain overall low numbers with an ultimate goal of total eradication. However, initial attempts to reduce the population by electro-fishing and netting were inefficient and destructive to non-targeted fish species. This was because fishing efforts were random and with a combined surface area of 74.7 square kilometres at full supply (Uytendaal 2003), random effort is hindered by the physical size of both lakes.

Carp can inflict major environmental and economic costs on both the private and public sectors by reducing water quality and degrading aquatic habitats (Koehn *et al* 2000). Carp have been implicated in macrophyte destruction, through direct grazing and physical uprooting of plants, increases in water turbidity and nutrient loads by continually resuspending sediments and reductions in invertebrate biomass and composition through predation. In addition to this, they compete with native and other desirable species for both food and space (Fletcher *et al* 1985. Brumley 1991. Koehn *et al* 2000 as cited in Walker and Freeman 2003).

The tendency for carp to aggregate for breeding and feeding was identified early in the Tasmanian program as a behavioural trait that could be exploited to quickly remove large numbers of fish. However, the location of carp

aggregations was only possible when carp created visual disturbances by moving into shallow water to spawn. Aggregations in deeper water were unpredictable and infrequently discovered by random fishing efforts.

Since 1997, the use of biotelemetry technology has enabled greater efficiency in the detection of carp movement and activity. The tendency for carp to aggregate during feeding and breeding episodes has been identified in other water bodies (Johnsen and Hasler 1977) and is the principle underlying the use of this technology. Male carp are first captured then measured and weighed to ascertain whether they are large enough to carry a radio-transmitter inside their gut cavity (transmitter \leq 2% body weight in air). If suitable, the carp are surgically implanted with radio transmitters and released back into Lake Crescent or Sorell. Their movements are tracked from a boat using a radio telemetry receiver attached to a directional antenna.

Transmitter-implanted carp or 'tracker fish' are regularly located and used as indicators of carp activity, movement and range. This enables IFS fishing operators to target fish aggregations through netting and electro-fishing with a higher degree of efficacy. In doing so, the impacts on non-target species such as the golden galaxid (*Galaxias auratus*), a species endemic to Lake Sorell and Crescent, are also reduced.

This paper will outline the equipment used and the procedures employed in the IFS Carp Management Program. Methodology for the surgical implantation of transmitters into carp is also outlined.

2.0 Telemetry Equipment

2.1 Transmitters

The IFS carp program uses internal ultrasonic transmitters, which are surgically implanted into the abdominal cavity of adult male carp. Initially the program attempted to source transmitters with long battery life and signals strong enough to transmit through water depths up to three metres in depth (maximum depth in Lake Sorell). Transmitters need to be small enough to be

implanted into the body cavity of mature male carp weighing greater than 1.5kg. It was decided that terrestrial transmitters would be suitable for use in Lakes Sorell and Crescent due to the shallow average depths across both lakes. The most suitable transmitters for these requirements were found to be terrestrial transmitters coated with epoxy resin and dipped in a layer of natural bee's wax or similar to ensure the unit would be completely waterproof. The first transmitters (model TX2-ICP-1) were supplied by Bio-Telemetry Tracking Australia (Biotel) with a 12 month battery life and a straight wire (whip) antenna. The transmitters emit radio frequencies between 150.7875 and 152.49375 MHz. These frequencies were chosen as they fall within permitted operating frequencies covered by a radio-communications class licence (low interference potential devices) under the *Commonwealth Radio Communications Act, 1992*. The signal emitted is a short "ping" with variable signal interval depending on the model of transmitter used. Most transmitters used have a pulse interval rate of approximately 55 cycles per minute but variations between 40-60 have been used.



Figure 1. Bio-Telemetry Transmitter with aerial.

Since the inception of the IFS carp biotelemetry program, different transmitter suppliers have been sourced and trialed. The largest manufacturer of transmitters for the Carp Management Program are supplied by Biotelemetry Tracking Australia (Biotel). Biotel transmitters are cylindrical, approximately 5 cm long with a diameter of 2 cm. They weigh 24 grams in air, are powered by lithium batteries, which last approximately one year and have a signal interval of 1.25 seconds. Because of the size of these transmitters, a minimum carp weight of approximately 1.5 kg is required for transmitter implant.

Other transmitters from various suppliers are constantly being sourced, trialed and evaluated. In general different transmitters are being trialed with a view to sourcing a transmitter that maximises battery life, durability, size, signal output and price. Other transmitter suppliers currently being utilised are outlined in table 1.

Manufacturer	Series model	Length (mm)	Diameter (mm)	Weight (grams in air)	Duty cycle	Battery life (months)
Bio-Telemetry Tracking Australia	(model TX2-ICP-1)	30	15	15	24hrs	4
Bio-Telemetry Tracking Australia	(model TX2-ICP-1)	50	20	24	24 hrs	12
Sirtrack Wildlife Tracking Systems	FRT conventional pulse transmitter	48	16	16.3	12hrs on / 12 hrs off	18
Sirtrack Wildlife Tracking Systems	FRT conventional pulse transmitter	40	14	10.3	12hrs on / 12 hrs off	9
Titely Electronics	F1850 prototype series	50	20	24	12hrs on / 12 hrs off	18
Titely Electronics	F1850 prototype series	20	12	11	12hrs on / 12 hrs off	9
ATS Advanced Telemetry Systems, INC	F1835 crystal controlled 2 stage series.	42	17	14	12hrs on / 12 hrs off	30
ATS Advanced Telemetry Systems, INC	F1855 crystal controlled 2 stage series.	68	17	25	12hrs on / 12 hrs off	44

Table.1 - Specifications for Transmitters used in Lakes Sorell and Crescent

Transmitter implant surgical procedures are based on those outlined by Johnsen and Hasler (1977). Unlike this study, the IFS carp program avoids holding fish captive for long periods both prior to, and after implant. This is due to an observed improvement in transmitter retention and fish survival when handling times are reduced as much as possible.

Since the inception of the biotelemetry program in 1997, between 10 and 15 active transmitters have been maintained in each lake. This range has proven adequate for detecting aggregations and scanning frequencies effectively at a moderate boat speed. When transmitter batteries approach their expiry date, implanted fish are retrieved. Their transmitters are then removed and the fish either destroyed or re-implanted with new transmitters. In some cases, if the transmitter is cleanly removed, the fish will be sewn up and placed in a holding pen for observation in the hope that it will fully recover and be available for future implantation. Expired transmitters are returned to the manufacturer for reprogramming and/or new batteries.

2.2 Telemetry Receiver Systems

Radio-transmitter implanted fish are individually identified by a unique radio frequency. The frequencies are programmed into a telemetry receiver and scanned continuously for approximately three seconds per frequency. This three second interval can be shortened or lengthened accordingly however the Carp Program has found a three second frequency interval most appropriate. When the receiver registers a programmed frequency, an audible "ping" sound will be heard through an in-built speaker system. The receiver can then be paused on that particular frequency so that an exact location of the tracker fish can be ascertained using a directional antenna. Therefore, in a typical tracking situation where 10 tracker fish are known to be functioning within a lake, it will take a telemetry receiver approximately 30 seconds to travel through an entire scan. This time period must be taken into account when choosing an appropriate boat speed to effectively radio track an area.

Each transmitter is programmed with a particular frequency ranging between 150.7875 and 152.49375 megahertz. Initially, two receivers, a Biotel RX3 and a Lotek Suretrack STR 1000 were both used to locate transmitter-implanted carp. The Biotel receiver has a manual search for a maximum of 9 pre-programmed frequencies.

This form of receiver made tracking a number of different frequencies from a boat difficult due to the need to physically change between frequencies while in motion. However, the Lotek receiver proved most suitable because it could be programmed to automatically search for up to 20 transmitters in each of up to four separate frequency (memory) partitions. Therefore, a theoretical limit of 80 transmitters is possible for use with the Lotek receiver. Separate partitions are used to store active transmitter frequencies in each lake.

The Lotek receiver scans the partition by sequentially sampling each frequency in the partition. The sample period for each frequency was set at between 2.5 to 3 seconds which is long enough to sweep a receiving aerial in a 180 degree arc and hear at least one of the “pings” which identifies a particular tracker fish in the vicinity. When a tracker frequency is located, the receiver can be locked onto that frequency so operators can move closer and accurately ascertain the tracker fish location. After location, the tracker frequency can be removed from the overall scan partition thereby shortening the overall scan time and increasing the efficiency of locating the other tracker fish.



Fig 2 . LOTEK Suretrack STR 1000 Telemetry Receiver.

2.3 Receiving Transmitter Signals

To receive signals, an antenna is attached to the LOTEK receiver using a flexible cable jack connector. The Carp Management Program uses a directional antenna or 'yagi'. The yagi antenna derives its name from one of the two Japanese inventors, Yagi and Uda, who developed the first directional antennae in the early 1920's. For radio-tracking on lakes such as Sorell or Crescent, the yagi is boat mounted using non-conducting PVC pipe as a mast when tracking. The yagi is mounted approximately 2 metres above the water surface for a number of reasons: adequate elevation can significantly increase signal strength, elevating the antenna can assist in dampening interference from outboard motors and also keeps the antenna out of the boat operators way. When tracker fish are located near to the lake-shore, a hand held yagi can be connected to the receiver and an exact location of the tracker fish can be found on foot.

Due to the directionality of the yagi, the strongest signals are received when the antenna is pointed directly towards the implanted fish. However, depending on the distance between the transmitter and receiver, an audible signal may be heard with the yagi pointed at more than 90 degrees from the direction of the transmitter. Because of the polarising effect on radio-waves of the water-air interface (Velle *et al* 1980), the strongest signal is received when the plane of the aerial is mounted perpendicular to the waters surface.




Fig 3. Radio tracking from boat using mast-mounted yagi.



Fig 4. Radio tracking an implanted carp using a hand held yagi

As illustrated in figures 4 and 5, the Carp Management Program uses 3 element, collapsible directional yagi antennas. Care must be taken to protect the elements of the antennas against bending and connection failure. Any

damage to an antenna can negatively effect frequency receiving strength and therefore tracking effectiveness.

3.0 Radio Tracking Techniques

Peak spawning times for carp have been identified as the warmer months from September through to March when water temperatures are higher (Drori 1994) During this period, the Carp Management Program monitor lakes Sorell and Crescent daily depending on weather conditions. Each time a lake is tracked, each transmitter fish that is located is given an approximate grid location. This is recorded on a prepared data sheet showing all operating tracker fish frequencies, their implant details and their last recorded position, (refer figure 5). Each data sheet is then entered into a computer database, which has been used to monitor the movements of individual male carp over time.




Fig 5. A biotelemetry field record data sheet for Lake Sorell, March 2003.

Between 10 and 15 tracker fish has proven to be a large enough sample of the population to detect aggregations in each lake. As the number of frequencies scanned increases, so does the chance of missing a signal due to passing the range of the signal before the frequency is scanned (10 frequencies takes approximately 30 seconds to scan). For larger frequency partitions, the boat speed and scan interval can be reduced to avoid these

problems. For each scan, the mounted yagi is manually rotated through a 180 degree arc. Because the yagi picks up frequencies approximately 90 degrees from its centre, this will effectively cover a 360 degree scan if direction of travel is constant over an entire frequency scan period.

The gain function of the receiver allows the operator to adjust the audibility of the signal and the level of background noise. Under normal circumstances, the Carp Program has found that a gain of around 80 dB set in the Lotek enables an audible signal to be detected over the background noise. With gain set at between 80 and 85 dB, a tracker fish can be detected up to 800m, and sometimes beyond 1200 metres, from the antenna depending on climatic conditions. As the distance to the transmitter is decreased, the gain can be reduced which improves the directionality of the yagi. This allows operators to determine the location of a tracker fish with a greater degree of accuracy. Experienced operators can, with very high precision, estimate *distance* to transmitter fish by the signal strength emitted at low gains. However, signal strength at a given gain can be influenced by several variables; water depth, turbidity and conductivity (Winter, 1979, Velle *et al* 1980) The height of the yagi and substrate topography will also effect the radio reception of transmitter frequencies travelling from water into air.

Diagram 1 illustrates how radio tracking can monitor the development of aggregations for immediate benefits and for future research and trend identification.




Diagram 1. Development of a carp aggregation in Lake Sorell, 2002

4.0 Surgical Implant Techniques

4.1 Introduction

Surgical implantation of radio transmitters into fish has been carried out on several species: chinook salmon, *Oncorhynchus tshawytscha* (Adams *et al* 1998); coho salmon, *Oncorhynchus kisutch* (Burger 1980); flathead catfish, *Pylodictis olivaris* (Hart and Summerfelt 1975); channel catfish, *Ictalurus punctatus* (Marty and Summerfelt 1986; Summerfelt & Mosier 1984); Colorado squawfish, *Ptychocheilus lucius* (Tyus *et al.* 1984); common carp, *Cyprinus carpio* (Johnsen and Hasler 1977; Berry 1982).

Accordingly, there is much literature describing surgical procedures of radio transmitter implantation. Despite this, several practical refinements to these surgical techniques have been developed during the long-term use of biotelemetry by the IFS Carp Management Program.

Most carp are implanted during colder months when water temperatures are low. This is to reduce infection rates after surgery and to ensure that an ample period of recovery time is available before carp attempt to aggregate for spawning purposes. Through 6 years of observation the carp team has concluded that implanted male carp return to 'normal' behaviour patterns in a relatively short space of time after successful implantation. The very fact that tracker fish are located in spawning and feeding aggregations only days after implantation supports these observations.

4.2 Surgical Equipment

General surgical equipment includes:

- stainless steel kidney dishes
- stainless steel scissors
- tweezers
- scalpels
- locking and needle nosed forceps
- suture needles
- cotton wool
- ethanol for sterilisation
- sterilised surgical gloves
- cotton gloves for holding the carp in place

All surgical operations were performed in a purpose-built mesh cradle designed to fit over a plastic fish bin. The cradle is a simple wooden frame

supporting a cloth mesh (figure 6). During surgery, the bin is filled with water and the cradle is positioned over the bin. The carp is held in an upside down position so that the surgical area is exposed but the head is emersed sufficiently to irrigate the gills. Observation has also shown that holding the carp in an upside down position seems to pacify the carp to some degree. Cotton gloves are used to hold the fish during the procedure.

A variety of sutures have been trialed, with varying success. Silk thread with pre-threaded needles proved to be adequate. However, a relatively inexpensive alternative is light (6 lb) mono-filament nylon fishing line. This material is strong, has good knot strength and is equally effective as silk. Dissolving sutures were also trialed but their use was aborted due to a high rate of suture failure. The suture needles determined to be most effective were size 6-8 flat point needles.

4.3 Sterilisation and fish handling procedures

Immediately prior to surgery, all surgical equipment is placed in a kidney dish and emersed in 100% ethanol. Excess ethanol is then decanted and the remaining ethanol and surgical tools “flamed” to remove residual ethanol and augment the sterilisation process. Heat sensitive equipment such as transmitters and sutures are emersed in ethanol and air-dried instead of flamed. In earlier implants, wounds were irrigated with iodine solution (Betadine) which is hypo-allergenic and non-toxic. However, the current consensus amongst operators is that wound irrigation is not necessary, as seepage from bleeding and mucous secretions appears to reduce infection.

Sterile practices are adopted in the field where practical. However, it appears that reducing stress is probably an equally important factor in reducing fish mortality, post-operative infection and the loss of transmitters through ruptured wounds. Keeping holding times to a minimum prior to, during, and after implant surgery is crucial in reducing stress. In earlier procedures, fish were captured and detained for up to a week in a holding pen and transported to shore based facilities for implant when required. Over time, it was

recognised that less handling time improved the recovery process and decreased mortality rates. The procedure has been refined to reduce handling, holding and procedural time, therefore reducing overall stress on the carp.

Also important in stress minimisation is ensuring that all equipment is ready for surgery and that the transmitter to be implanted is operational. In addition, the transmitter should be tuned in to the receiver and the serial number recorded to ensure that the fish frequency is recorded correctly.

4.4 Anaesthesia

'AquiS' is a solution consisting of clove oil and is commonly used as a fish anaesthetic. Prior to surgery, carp are placed in a 20 litre tub of fresh water which has been diluted with AquiS at a rate of 0.3 mg/l. The carp is observed until it is unable to right itself for any period of time and its swimming function is lost. At this point the carp is removed from the anaesthetic and placed on a cradle partly submerged in fresh water ready for surgery. After surgery the carp is placed into a 20 litre tub of aerated fresh water and kept under observation until fully recovered and ready for release.

4.5 Surgical Procedure

Fish chosen as implant recipients are large mature males. Carp of 1.5 kg are deemed to be adequate, however fish greater than 2.0 kg are preferable. This is because for successful implantation, it is recommended that fish should not be implanted if the weight of the transmitter is greater than 2% of the actual body weight of the fish when weighed in air (Winter 1979). Prior to implant surgery, the sex of fish is ascertained by gentle hand stripping. If a fish does not express milt but appears to be a male, the sex is checked visually after the body cavity is opened and gonad tissue can be observed. Accordingly, if this process reveals female gonads, the operation is aborted and the fish is despatched and processed.

The surgical process requires at least two personnel: one to perform the surgery and another to hold the fish in an inverted position in the cradle. The latter person secures the fish with both gloved hands; one hand around the tail and the other placed around the pectoral girdle whilst ensuring that the gill opercula are not held shut.

When the fish is firmly secured, the incision site is prepared by removing scales along a 50-70 mm line (dependant on size of transmitter) slightly offset on either side of the mid-ventral line, posterior to the pelvic fin and anterior to the anus (see figure 6). Incisions along the mid-ventral line are possible but the risks of cutting the gut or a major blood vessel are high. Abrasion with the substrate and possible infection can also occur if the incision is made at the mid-ventral line.

With a scalpel, the first incision should be shallow without penetrating the inside of the body wall. The length of the cut should be between 30 to 40 mm and just large enough to permit transmitter insertion (refer figure 7).

Using tweezers and grasping one side of the incision, lift the wall away from the gut. This reduces the chance of cutting visceral organs or gonad. Complete the incision through the body wall whilst exercising caution not to sever any viscera. At this stage, the sex of the fish and the stage of gonadal maturation can be established (if uncertain). Under normal circumstances, there will be a small to moderate amount of bleeding. This may be beneficial by irrigating the wound, thus expelling pathogens and promoting the healing process.



Figure 6 Removing scales and making the first incision

Holding the fish firmly with the transmitter aerial trailing posteriorly, push the transmitter into the body cavity. Initially, angle the blunt end down then gently push the end towards the head angling the leading edge outward toward the body wall to avoid disturbing the organs (refer figure 8). Continue until the transmitter lies neatly within the body cavity. To secure the transmitter in place, the antenna is then threaded on to a suture needle and is sewn through the gut wall, exiting through the body wall on the posterior end of the incision. Alternatively, needle nosed forceps can be used to push the aerial through the gut wall with the aid of a very small incision made with a scalpel.



Figure 7 Implanting the transmitter into body cavity



Figure 8 Transmitter being positioned in body cavity

Several 200 mm lengths of mono-filament should be prepared and pre-threaded onto suture needles. Using locking forceps to hold the suture needle, each suture is started about 5 mm on either side of the wound. The needle is passed through the skin and through the muscle, exiting from the side of the incision at about half the thickness of the body wall. The incision is then closed by entering the muscle on the other side of the wound and again, at about half the thickness of the body wall, then out through the skin about 5 mm from the edge of the incision. The suture is then tied off using a standard locking knot.



Figure 9. The implanted transmitter securely sutured and ready for release.

The sutures are repeated about 5-7 mm apart until the wound is sutured completely shut. The tails of each suture should be trimmed back to about 2 mm from the knot. To further identify fish, number-coded T-bar tags that are implanted beside the dorsal fin of each fish. This is useful for two reasons: firstly, fish can be individually identified when recaptured without using the receiver and secondly; if the transmitter is “dropped” and the fish recaptured at a later date, the fish can still be identified.

5.0 Discussion

Currently, there are carp eradication and research programs operating in all states of Australia. However in most cases other than Tasmania, biotelemetry is used as a means to monitor movement, range and growth of a carp population (Stuart and Jones 2002). The Tasmanian Carp Management Program is unique in that it uses biotelemetry as a primary tool in the removal of carp during spawning and feeding aggregations. Since the Carp Management Program began using biotelemetry in 1997, a total of 4971 and 1986 have been removed from lakes Crescent and Sorell respectively. The Carp Program estimates that over 90% of these carp have been captured and removed as a result of radio tracking and location operations.

Radio tracking has also identified certain trends and key habitat areas, which have been frequented by different tracker fish over long periods of time. This information has allowed the carp program to monitor these areas closely and in some cases to build static carp traps.

These traps have captured carp with no direct fishing effort required. In addition to this, two key areas of habitat in Lake Sorell have been fenced off due to repeated spawning aggregation attempts. In this case biotelemetry has greatly assisted in identifying these areas. The fencing off of these areas will force future carp aggregations into marginal habitat areas where spawning

success will be reduced and aggregations will be more vulnerable to fishing efforts.

The ability to use radio tracking to locate key aggregations of carp and remove a large percentage in one fishing session, has increased fishing efficiency and lowered catch per unit effort markedly. The chance to remove large numbers of carp during these events allows the Carp Management Program team to maximise the use of equipment and personnel at pivotal times rather than trying to spread these resources randomly across an entire season. There certainly exists ambiguity in relation to which fishing efforts are directly related to radio tracking location events or indirectly related to radio tracking trends and patterns observed over time. To adequately illustrate this, it would be useful to examine the direct relationship biotelemetry has on catch effort and fishing efficiency more closely in a separate document.

As the population of carp becomes lower in lakes Crescent and Sorell, random fishing techniques over such a large surface area obviously become much less effective. For this reason, the Carp Management Program will rely on biotelemetry more heavily as the population of carp is further reduced. Furthermore, as the population of carp decreases in lakes Crescent and Sorell, the availability of suitable male carp for transmitter implant will also be challenged. This is one reason why different transmitters are being trialed in the biotelemetry program in an attempt to find suitable equipment and techniques to implant either smaller, sexually determined carp, or have presently implanted carp last as viable tracker fish for longer periods of time.

6.0 References

Adams N.S., Rondorf W., et al. (1998). "Effects of Surgically and Gastrically Implanted Radio Transmitters on Growth and Feeding Behaviour of Juvenile Chinook Salmon." Transactions of the American Fisheries Society 127: 128-136.

Adams N.S., Rondorf W., et al. (1998). "Effects of surgically and gastrically implanted radio transmitters on swimming performance and predator avoidance of juvenile chinook salmon (*Oncorhynchus tshawytscha*)." Canadian Journal of Fisheries and Aquatic Science 55: 781-787.

Armstrong J. D., Lucas M., et al. (1988). "A combined radio and acoustic transmitter for fixing direction and range of freshwater fish (RAFIX)." Journal of Fish Biology 33: 879-884.

Armstrong J. D., Herbert N.A., (1997). "Homing movements of displaced stream-dwelling brown trout." Journal of Fish Biology 50: 445-449.

Baras E., Westerloppe L. (1999). "Transintestinal Expulsion of Surgically Implanted Tags by African Catfish *Heterobranchus longifilis* of Variable Size and Age." Transactions of the American Fisheries Society 128: 737-746.

Burger C. (1980). radio tracking of Adult Coho Salmon (*Oncorhynchus kisutch*) in the Deshka River, Alaska. Anchorage, U.S fish and Wildlife Service.: 151-158.

Close T. L. (2000). "Detection and Retention of Postocular Visible Implant Elastomer in Fingerling Rainbow Trout." North American Journal of Fisheries Management 20: 542-545.

Cohen M.A. (1998). "A low-budget and versatile radio tracking system." Fisheries Management and Ecology 5: 491-500.

Drori S., Ofir M., et al. (1994). "Spawning Induction in common carp (*Cyprinus carpio*) using pituitary extract or GNRH superactive analogue combined with metoclopramide - analysis of hormone profile, progress of oocyte maturation and dependence on temperature." *Aquaculture* 119(4): 393-407.

Gonczi A. P. "Telemetry experiments with newly stocked brown trout (*Salmo trutta* L) in river reservoirs in Northern Sweden". Harnosand, National Board of Fisheries: 212-216.

Guy C. S., Willis W., et al. (1994). "Biotelemetry of White Crappies in a South Dakota Glacial Lake." *Transactions of the American Fisheries Society* 123: 63-70.

Hart L. G., Summerfelt R.C. (1975). "Surgical Procedures for Implanting Ultrasonic Transmitters into Flathead catfish (*Pylodictis olivaris*)."
Transactions of the American Fisheries Society 104: 56-69.

Houston A.H., Czerwinski C.L., et al. (1973). "Cardiovascular-respiratory activity during recovery from anesthesia and surgery in brook trout (*Salvelinus fontinalis*) and carp (*Cyprinus carpio*)."
Journal Fisheries Board Of Canada 30(11): 1705-1712.

Johnsen P., Hasler A. (1977). "Winter Aggregations of Carp (*Cyprinus carpio*) as Revealed by Ultrasonic Tracking." *Transactions of the American Fisheries Society* 106(6): 556-559.

Knights B., Lasee B. (1996). "Effects of Implanted Transmitters on Adult Bluegills at Two Temperatures." *Transactions of the American Fisheries Society* 125: 440-449.

Koehn, J., Brumley A.R., et al. (2000). *Managing the Impacts of Carp*. Canberra Bureau of Rural Sciences.

Marty G., Summerfelt R.C. (1986). "Pathways and Mechanisms for Expulsion of Surgically Implanted Dummy Transmitters from Channel Catfish." Transactions of the American Fisheries Society 115: 577-589.

Mulford C.J. (1982). "Use of a surgical skin stapler to quickly close incisions on striped bass." Management Briefs: 571-573.

Petering R., W., Johnson D.L. (1991). "Suitability of a cyanoacrylate Adhesive to close Incisions in Black Crappies Used in Telemetry Studies." Transactions of the American Fisheries Society 120: 535-537.

Roberts J., Chick A., et al. (1995). "Effect of Carp, *Cyprinus carpio* L, an Exotic Benthivorous Fish, on Aquatic Plants and Water Quality in Experimental Ponds." Marine and Freshwater Research 46: 1171-1180.

Stuart I., Jones M. (2002). Ecology and Management of common carp in the Barmah-Millewa forest. Melbourne, Arthur Rylah Institute for Environmental Research.

Summerfelt R.C., Smith L. S. (1979). Anesthesia, Surgery and Related Techniques. A Handbook On Biotelemetry and Radio Tracking. Amlaner C.J., Macdonald D.W, eds. New York, Pergamon: 213-272.

Summerfelt R.C., Mosier D. (1984). "Transintestinal Expulsion of Surgically Implanted Dummy Transmitters by Channel Catfish." Transactions of the American Fisheries Society 113: 760-766.

Tyus H.M., Burdick B., et al. (1984). "Use of radiotelemetry for Obtaining Habitat Preference Data on Colorado Squawfish." North American Journal of Fisheries Management 4: 177-180.

Uytendaal Adam. (2003). "Water quality in lakes Sorell and Crescent: Underlying processes and management options". Rehabilitation of lakes

Sorell and Crescent Report Series no. 5. Inland Fisheries Service, Tasmania. 7-8.

Velle J. I., Lindsay J.E., et al. (1980). "An Investigation of the Loss Mechanisms Encountered In Propagation From A Submerged Fish Telemetry Transmitter". Paper of Ph.D. Thesis, University of Wyoming, Fall 1979.

Venditti D.A., Rondorf D. W., et al. (2000). "Migratory Behavior and Forebay Delay of Radio-Tagged Juvenile Fall Chinook Salmon in a Lower Snake River Impoundment." *North American Journal of Fisheries Management* 20: 41-52.

Wagner G.N., Stevens D. E., et al. (2000). "Effects of Suture Type and Patterns on Surgical Wound Healing in Rainbow Trout." *Transactions of the American Fisheries Society* 129: 1196-1205.

Walker R., Freeman R. (2003). "Lakes Sorell and Crescent Fisheries Management. Integrated and multi-disciplinary approach to the rehabilitation of Lakes Sorell and Crescent" Inland Fisheries Service Tasmania. Page 1.

Walsh M.G., Bjorgo K. A., et al. (2000). "Effects of Implantation method and Temperature on Mortality and Loss of Simulated Transmitters in Hybrid Striped Bass." *Transactions of the American Fisheries Society* 129: 539-544.

Winter, J. (1979). *Underwater Biotelemetry. A Handbook on Biotelemetry and Radio Tracking*. C. J. Amlaner and D. W. Macdonald. New York, Pergamon Press: 371-395.